

Rothamsted Repository Download

A - Papers appearing in refereed journals

Rottensteiner, H. and Theodoulou, F. L. 2006. The ins and outs of peroxisomes: co-ordination of membrane transport and peroxisomal metabolism. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*. 1763 (12), pp. 1527-1540.

The publisher's version can be accessed at:

- <https://dx.doi.org/10.1016/j.bbamcr.2006.08.012>

The output can be accessed at: <https://repository.rothamsted.ac.uk/item/89qzx/the-ins-and-outs-of-peroxisomes-co-ordination-of-membrane-transport-and-peroxisomal-metabolism>.

© 2006, Elsevier

Review

The ins and outs of peroxisomes: Co-ordination of membrane transport and peroxisomal metabolism

Hanspeter Rottensteiner^{a,*}, Frederica L. Theodoulou^b

^a Medical Faculty of the Ruhr-University of Bochum, Department of Physiological Chemistry, Section of Systems Biochemistry, 44780 Bochum, Germany

^b Crop Performance and Improvement Division, Rothamsted Research, Harpenden, AL5 2JQ, UK

Received 23 June 2006; received in revised form 15 August 2006; accepted 18 August 2006

Available online 23 August 2006

Abstract

Peroxisomes perform a range of metabolic functions which require the movement of substrates, co-substrates, cofactors and metabolites across the peroxisomal membrane. In this review, we discuss the evidence for and against specific transport systems involved in peroxisomal metabolism and how these operate to co-ordinate biochemical reactions within the peroxisome with those in other compartments of the cell.

© 2006 Elsevier B.V. All rights reserved.

Keywords: ABC transporter; β -oxidation; Adenine nucleotide translocator; Redox shuttle; Peroxisomal pH; Porin

1. Introduction

Peroxisomes are near-ubiquitous organelles of eukaryotic cells which perform a range of functions, many of which are organism- or tissue-specific [1,2]. The common metabolic activities of peroxisomes are the oxidation of fatty acids and generation and removal of hydrogen peroxide, but peroxisomes have also been implicated in β -oxidation of aromatic and cyclic compounds, synthesis of plasmalogens, isoprenoids, penicillin, lysine and glycine betaine, metabolism of purines and pyrimidines, and catabolism of polyamines, D-amino acids and methanol. Dependent on the organism, these organelles also participate in the glyoxylate cycle, pentose phosphate pathway and photorespiration and key roles for peroxisomes in signalling and development have emerged recently ([3,4], and references therein). This diversity of function is often reflected in peroxisomal specialisation, especially in plants [5] and also in trypanosomes [6,7].

The multiple roles of peroxisomes imply not only that a relatively large number of chemically diverse solutes must enter the peroxisome, but also that the products of peroxisomal metabolism must be exported from the organelle. Moreover,

since a number of pathways are shared between the peroxisome and other cellular compartments, the metabolic steps that occur within the peroxisomal matrix or on the inner face of the peroxisomal membrane must be integrated with those in other cellular compartments to permit efficient operation of biochemical pathways.

The transport properties of the peroxisomal membrane have been a controversial issue for several decades. Early studies with isolated organelles revealed a lack of structure-linked latency for certain peroxisomal enzymes, suggesting that these organelles are freely-permeable to substrates (for more details see Antonenkov and Hiltunen, this issue). Although critics of this hypothesis have argued that the *in vitro* permeability of the membrane reflects damage and disruption during isolation (e.g. [1]), there have been a number of reports of relatively non-specific porin-like proteins in the peroxisomes of mammals, plants and yeast (reviewed in [8]). Recent, rigorous studies with isolated peroxisomes indicate that these organelles are permeable to low molecular weight, hydrophilic solutes such as glycolate and urate, but not to more bulky cofactors [9]. The latter conclusion is strongly supported by genetic studies which demonstrate that the peroxisomal membrane of *Saccharomyces cerevisiae* is impermeable to NAD(H) and acetyl-CoA *in vivo* [10]. Although the functions of many peroxisomal membrane proteins remain unknown, a growing number of peroxisomal

* Corresponding author.

E-mail address: Hanspeter.Rottensteiner@rub.de (H. Rottensteiner).

transport proteins have been characterised at the molecular level and these form the subject of this review. Of these, the best characterised are the adenine nucleotide translocator and ATP-binding cassette (ABC) transporters which have been implicated in transport of substrates for β -oxidation.

2. Import of substrates, co-substrates, and cofactors for β -oxidation

2.1. Import of substrates for β -oxidation: the role of ABC transporters

β -Oxidation not only provides a major pathway for the degradation of fatty acids but also serves a synthetic role in the production of a range of bioactive, lipid-derived molecules such as jasmonates, plasmalogens and bile acids [1,11]. In yeast and plants, the peroxisome is the sole site of fatty acid β -oxidation, whereas in mammals, very long chain fatty acids (VLCFA) are first chain-shortened in the peroxisome prior to complete oxidation in mitochondria. Oxidation of long chain fatty acids (LCFA) is predominantly mitochondrial and short- and medium chain fatty acids (S- and MCFA) are oxidised exclusively in mitochondria. In all cases, formation of fatty acyl-CoA esters is a prerequisite for entry into the β -oxidation spiral and is catalysed by members of a family of acyl-CoA synthetase enzymes, also known as activases, which differ in their substrate specificity. The intracellular location of fatty acid activation differs according to chain length and organism and thus determines the identity of the molecular species which must traverse the peroxisomal membrane. SCFA and MCFA are considered to be readily membrane-permeable and are able to enter mitochondria and peroxisomes in their protonated form. Esterification of free fatty acid to CoA, however, generates an amphipathic molecule, which is unable to cross membranes in the absence of a transport protein. Genetic studies have provided strong evidence for the involvement of members of the ABC protein superfamily in the peroxisomal import of substrates for β -oxidation [3].

A functional ABC protein comprises two copies each of two units: the membrane spanning domain (MSD), also known as the transmembrane domain (TMD) and the soluble nucleotide binding fold (NBF) [12]. In eukaryotes, these units are frequently contained within a single polypeptide with the topology TMD-NBF-TMD-NBF, an arrangement commonly referred to as a “full size” transporter. Alternatively, two “half size” transporters (TMD-NBF) may dimerise to form a functional ABC protein. Peroxisomal ABC transporters belong to subclass D, almost all members of which are half size transporters. The ABCD subfamily contains between one and five members and is represented in all eukaryotic taxa studied to date [3,13–22], but has been most extensively studied in mammals, baker’s yeast and the model plant, *Arabidopsis thaliana*.

2.1.1. Yeast

Saccharomyces cerevisiae contains two peroxisomal ABC transporters, Pxa1p and Pxa2p (also known as Pat2p/Pal1p and

Pat1p), which have been shown by genetic and co-immunoprecipitation studies to form a functional heterodimer in the peroxisomal membrane [23–26]. *pxa1* Δ and *pxa2* Δ deletion mutants are unable to grow on the LCFA oleate as the sole carbon source, but growth on MCFA such as laureate was unaffected, suggesting that these transporters play a specific role in β -oxidation of LCFA. This agrees with the finding that MCFA are imported into the peroxisome as free fatty acids where they are activated by the acyl-CoA synthetase Faa2p [23]. In contrast, LCFA are activated in the cytosol and LCFA-CoA esters must be transported across the peroxisomal membrane (Fig. 1). Biochemical studies with permeabilised spheroplasts provided evidence that Pxa2p is required for import of LCFA-CoA but not of SCFA-CoA [27] and mistargeting of Faa2p to the cytosol was shown to make β -oxidation of MCFA dependent on Pxa1p/Pxa2p [23], providing further support that this is an acyl-CoA transporter [28].

2.1.2. Plants

The genome of *Arabidopsis thaliana* contains a single ABCD gene, known as COMATOSE (CTS) and also as PXA1, PED3 and *AtABCD1* [15,29,30]. Interestingly, unlike non-plant ABCD proteins, CTS is a full-length ABC transporter, in which two dissimilar half-size units are contained in a single polypeptide. All plant ABCD proteins identified to date have this modular architecture, suggesting that the fusion of two halves was a single, ancient event in evolutionary history. CTS is expressed ubiquitously in *Arabidopsis* and, unsurprisingly, null alleles have a pleiotropic phenotype: *cts* mutants fail to germinate in the absence of classical dormancy-breaking treatments and seedlings require sucrose for establishment [15,31]. Oil bodies and fatty acyl-CoAs, particularly VLCFA-CoAs, are retained in mutant seedlings, indicating an inability to break down storage lipid [15]. It has been suggested therefore, that CTS either transports or regulates the import of fatty acids or fatty acyl-CoAs into the peroxisome [3,11,15,29,30]. The transported species is still a matter for debate: whilst the retention of VLCFA-CoA is suggestive that CTS transports acyl-CoA, it is possible that the CoA pool represents a sink for free fatty acids and thus that CoA retention does not reflect cytosolic accumulation of CTS substrates. Moreover, the characterisation of a double peroxisomal acyl-CoA synthetase mutant provides evidence that free fatty acids could be transported into plant peroxisomes. *Arabidopsis* has two peroxisomal acyl-CoA synthetases (LACS 6 and 7) which have a substrate preference for (V)LFCA [32]. The active sites of LACS 6 and 7 are within the peroxisome and the *lacs6*, *lacs7* double mutant is unable to degrade storage lipid and cannot establish in the absence of sucrose [32,33]. Furthermore, peroxisome morphology of β -oxidation mutants such as *kat2* (thiolase) and *mfp2* (multifunctional protein) is distorted due to intraperoxisomal accumulation of acyl-CoAs [34,35], whereas it is normal in *cts* and *lacs6*, *lacs7*, suggesting that knocking out either the CTS transporter or the peroxisomal acyl-CoA synthetases prevents accumulation of acyl-CoAs in peroxisomes [36].




Fig. 1. Transport processes associated with β -oxidation. Dependent on chain length and organism, fatty acids are imported into peroxisomes as free acids (FFA) or CoA esters (FA-CoA). CoA esters formed in the cytosol or on the endomembrane system by the action of acyl-CoA synthetases (ACS) are transported across the peroxisomal membrane by one or more ABC proteins, at the expense of ATP hydrolysis. Alternatively, short- and medium chain free acids may enter the peroxisome by passive transport and are activated by peroxisomal ACS. This reaction requires intraperoxisomal pools of ATP and CoA which are supplied by the peroxisomal adenine nucleotide translocator (ANT) and by an unknown mechanism (?) respectively. AMP and pyrophosphate (PP_i) are produced as by-products of fatty acid activation; AMP is exchanged for ATP by ANT and pyrophosphate is thought to be converted to inorganic phosphate which is exported by an NEM-insensitive phosphate transport protein (PT).

Other aspects of the *cts* mutant phenotype suggest that CTS could be a transporter of broad substrate specificity. Roots of *cts* mutants are resistant to indole butyric acid (IBA) and 2,4 dichlorophenoxybutyric acid (2,4-DB), indicating that they are unable to metabolise these compounds by β -oxidation to the corresponding active auxins [15,29,30]. The simplest explanation for these data is that CTS transports IBA and 2,4-DB or their CoA esters. Both basal and wound-induced jasmonic acid (JA) levels are reduced in *cts* mutants, suggesting that CTS is also involved in peroxisomal import of jasmonate precursors which yield JA following reduction by the peroxisomal enzyme OPR3 and three rounds of β -oxidation [37]. Many JA biosynthetic and signalling mutants are male sterile but *cts* mutants apparently have sufficient JA to retain fertility, suggesting an additional, probably passive, route for import of JA precursors. Nevertheless, fertilisation is reduced in *cts* mutants, which exhibit defects in pollen tube growth. This phenotype is not rescued by application of JA, suggesting an additional role for CTS in optimal fertility in *Arabidopsis* (S. Footitt, D. Dietrich, A. Baker, M. Holdsworth and F. Theodoulou, unpublished data). Taken together, comparisons of the phenotypes of *cts* mutants with those defective in other steps of β -oxidation is pointing to a role of β -oxidation not only in energy metabolism but also in the generation or processing of developmental signals and new roles for peroxisomal metabolism may await discovery in plants. An exciting challenge for the future will be the determination of the precise biochemical role of CTS, which awaits transport studies.

2.1.3. Mammals

The ABCD subfamily of humans and mice has four members: Adrenoleukodystrophy protein (ALDP; ABCD1), Adrenoleukodystrophy related protein (ALDR; ABCD2), PMP70 (ABCD3) and PMP69 (P70R; ABCD4), which exhibit overlapping but distinct tissue expression patterns [38–43]. These proteins have been shown to be integral peroxisomal membrane proteins [40,41,43–46] and a search for interacting partners revealed that ALDP, ALDR and PMP70 interact with PEX19, a cytosolic acceptor for peroxisomal membrane proteins which is involved in their intracellular sorting and trafficking to the peroxisomal membrane [47–50].

PMP70 was the first peroxisomal ABC transporter to be identified and characterised functionally and is expressed predominantly in liver [40], but transcripts have been detected in all tissues investigated [41]. Overexpression of rat PMP70 in Chinese hamster ovary cells led to an increase in LCFA but not VLCFA β -oxidation by peroxisomes and site-directed mutagenesis demonstrated a requirement for conserved motifs in ATP binding and hydrolysis [51]. Unlike many genes encoding peroxisomal proteins, mutations in *ABCD3* are not associated with disease in humans and the physiological significance of PMP70 awaits investigation [52,53]. Preliminary reports from *abcd3*^{-/-} knockout mice suggest an additional role in transport of 2-methylacyl-CoA esters [54]. On the basis of sequence homology, PMP69 has been suggested to play a similar role to PMP70, though this transporter has not yet been characterised functionally [38].

ALDP has received much attention due to its role in the X-linked neurodegenerative disease, Adrenoleukodystrophy (X-ALD), which is characterised biochemically by the accumulation of VLCFA in plasma and tissue. Although X-ALD was originally associated with reduced activity of VLC acyl-CoA synthetase [55–57], positional cloning identified the *ALD* gene as an ABC transporter [42] and transfection with the wild type *ABCD1* cDNA was shown to correct the VLCFA β -oxidation defect of X-ALD fibroblasts [58–60]. Currently, over 650 different mutations in the *ABCD1* gene have been reported [52; www.X-ALD.nl]; in the majority of cases investigated, immunoreactive ALD protein is absent, suggesting the existence of an efficient retrieval mechanism for mutant forms of ALDP or that non-functional ALDP is unstable [46,61,62]. The precise relationship between the clinical symptoms of X-ALD, the biochemical phenotype and mutated *ABCD1* is not currently clear [63,64]; for example, mouse models have been developed for X-ALD and although *abcd1*^{-/-} mice replicate the biochemical defects of patients, they do not exhibit the clinical and pathological symptoms until very late in life [65–67]. It has been proposed that elevated VLCFA levels in X-ALD are related to mitochondrial dysfunction [68], although this has been contested by a detailed ultrastructural study of mitochondria [69]. Enhanced microsomal fatty acid chain elongation has also been suggested as an additional mechanism underpinning VLCFA accumulation in X-ALD [70]. Despite the uncertainty regarding the pathology of X-ALD, it remains plausible that ALDP is a VLCFA or VLCFA-CoA transporter [52], and support for this hypothesis has been obtained by substrate-induced conformational changes [71], although unequivocal proof awaits direct demonstration of transport activity for purified ALDP.

Complementation studies have shown that expression of ALDRP and, surprisingly, PMP70 restores VLCFA β -oxidation in X-ALD fibroblasts, preventing VLCFA accumulation [58,63,72–74]. Pharmacological induction of endogenous peroxisomal transporters also resulted in restoration of VLCFA β -oxidation, providing promising leads for therapy [73,74]. These data suggest that mammalian peroxisomal ABC transporters are at least partially redundant and functional overlap between ALDP and ALDR has been confirmed by studies with knockout mice [75].

It has been suggested by several authors that the four half size transporters of the mammalian ABCD subfamily might heterodimerise to yield variants with modified transport properties [39,43,76], by analogy to human transporters associated with antigen processing (TAP) belonging to ABC subfamily B [77]. A number of studies have provided evidence for both homo- and heterodimerisation of ALDP, ALDR and PMP70 *in vitro* [63,78,79]. However, the expression patterns imply that, in several cell types, heterodimerisation is not physically possible *in vivo* [41,80,81] and this view is supported by the differential induction by pharmacological agents [82]. Moreover, purification of native ALDP and PMP70 strongly suggest that homomeric interactions predominate *in vivo* [83]. Nevertheless, heterodimerisation cannot be ruled out under specific conditions and is an important consideration for X-ALD gene therapy.

2.2. Transport of adenine nucleotides

The peroxisomal transporter best characterised is the adenine nucleotide transporter of *S. cerevisiae*, which is required for the activation of MCFA, the only established ATP-consuming reaction inside peroxisomes (Fig. 1). As discussed above, while the cytosolic fatty acid activation enzymes preferentially use LCFA as substrates, peroxisomes are the predominant site for MCFA activation [23]. The latter reaction is dependent on the supply of intraperoxisomal ATP and is catalysed by peroxisomal acyl-CoA synthetase *Faa2p*. Upon formation of the fatty acyl-CoA product, AMP is generated, which has to be exported back to the cytosol due to the lack of an adenylate kinase or an ATP regeneration system within peroxisomes. This cross-membrane traffic of adenine nucleotides (ATP in, AMP out) is catalysed by *Ant1p* [84,85], a member of the mitochondrial carrier family (MCF) generally involved in solute transport [86]. Both genetic and biochemical evidence supports this view: cells mutated in either *FAA2* or *ANT1* grow normally on LCFA but fail to do so on MCFA. In combination with a disabled *Pxa1p*–*Pxa2p* transporter, the *faa2* Δ and the *ant1* Δ mutants are unable to metabolise any fatty acid [23,84,85]. The β -oxidation rates of MCFA of an *ant1* Δ strain are also reduced when measured with intact cells. However, the respective cell lysates show a normal level of β -oxidation, which is consistent with a blockage in the ATP transport across the peroxisomal membrane in intact *ant1* Δ cells [85]. Likewise, heterologous expression of a peroxisomally targeted luciferase, which emits light in an ATP-dependent manner, required *Ant1p* for activity, but not when assayed in cell lysates [85].

Functional reconstitution of purified *Ant1p* into liposomes revealed the ability of *Ant1p* to transport adenine nucleotides specifically. ATP uptake was measured with proteoliposomes loaded with ATP, ADP, AMP, or the respective desoxynucleotides, whereas uptake was very low with other nucleotides or a number of other substrates for various known transporters [84]. The established inhibitors of the mitochondrial ADP/ATP carrier, carboxyatractyloside and bongkreic acid [87] did not reduce the activity of *Ant1p*, indicating that *Ant1p* represents a novel type of MCF transporter. Interestingly, *Ant1p* exhibits two transport modes, not only the exchange described above, but also the uniport of adenine nucleotides [88]. *In vivo*, the former mode serves to exchange ATP for AMP under ongoing β -oxidation, whereas the latter is probably most important for loading peroxisomes with ATP early in their genesis (Fig. 1). Since both modes are proton-compensated and electroneutral [88], import of negatively charged ATP should contribute to the formation of a proton gradient across the peroxisomal membrane. Consistent with this idea, the Δ pH across the peroxisomal membrane is significantly diminished in an *ant1* Δ knock-out strain as measured by pH-sensitive variants of fluorescent proteins [88,89]. The significance of such measurements will be discussed below.

Pmp47p from *Candida boidinii* represents a functional homolog of *Ant1p*. Sakai and colleagues showed that a *pmp47* Δ mutant also exhibits a specific growth defect on MCFA, which can be rescued by introducing *S. cerevisiae* *Faa2p*

devoid of its peroxisomal targeting signal [90]. Growth on MCFA was also restored upon expression of cytosolic Faa2p in *S. cerevisiae ant1Δ* cells [85] providing evidence that Pmp47p and Ant1p function as peroxisomal ATP carriers. Given that MCFA are now activated by truncated Faa2p in the cytosol and transported into peroxisomes through Pxa1p–Pxa2p, intraperoxisomal ATP becomes dispensable. That heterologous expression of PMP47 is able to restore growth of the *S. cerevisiae ant1Δ* mutant on MCFA [84] lends further credence to the conception that Pmp47p represents the *C. boidinii* ortholog of Ant1p.

The human MCF member PMP34 localises to peroxisomes and its sequence is closely related to Pmp47p and Ant1p [91]. PMP34 likely represents the mammalian ortholog of Ant1p, as it was able to partially complement the MCFA β -oxidation defect of the *ant1Δ* mutant, and showed some ATP exchange activity when reconstituted into liposomes [92]. However, due to the current lack of reports on the effects of PMP34 knock-down by RNAi or on PMP34 knock-out mice, its physiological role remains elusive.

A proteomic approach has led to the identification of a candidate peroxisomal adenine nucleotide translocator from *Arabidopsis*. PMP38 exhibits 25% and 26% amino acid identity to human PMP34 and *C. boidinii* PMP47, respectively and was shown by a combination of cell fractionation and immunocytochemical analysis to be an integral peroxisomal membrane protein [93]. Although the pattern of developmental changes in PMP38 abundance strongly suggest involvement in β -oxidation, the *in vivo* function has yet to be determined, for example, by characterisation of knock-out mutants or antisense lines.

2.3. CoA and phosphate transport

Peroxisomal acyl-CoA synthetases such as yeast Faa2p as well as 3-ketoacyl-CoA thiolases require free CoA within peroxisomes. Although some CoA probably enters the peroxisomal matrix thioesterified with LCFA via the Pxa1p–Pxa2p transporter of yeast, stoichiometry suggests that a certain pool of CoA must be provided by other means to maintain MCFA activation and the flux through the β -oxidation spiral. Furthermore, activation and degradation of MCFA occur normally in *pxa1Δpxa2Δ* mutant cells [23]. It is currently unclear whether peroxisomes contain a CoA transporter as has been suggested for mitochondria. Here, the MCF member Leu5p is required for the accumulation of CoA in the mitochondrial matrix and the related human Grave's disease protein can functionally replace Leu5p [94]. It is not entirely impossible that this carrier also localises to peroxisomes.

Intraperoxisomal ATP-dependent fatty acid activation generates pyrophosphate that probably decomposes into two molecules of orthophosphate, which have to leave the peroxisome for regeneration of ATP. Evidence for a phosphate transporter was recently provided using proteoliposomes reconstituted from peroxisomal membranes that had been purified from bovine kidney [95]. Although the molecular identity of this transporter remains to be determined, it differed from the mitochondrial transporter in terms of sensitivity

towards N-ethylmaleimide (NEM) and p-hydroxyphenylglyoxal (PHGP), as only the mitochondrial activity was adversely affected by these compounds [95].

2.4. Import of cofactors and regeneration of reducing equivalents

Genetic studies and transport assays utilising isolated peroxisomes suggest that the peroxisomal membrane is not freely permeable to bulky cofactors, implying that the peroxisome has separate pools of NAD^+ , NADP^+ , and FAD [9,10]. Acyl-CoA oxidase absolutely requires FAD, but the holoenzyme is assembled in the cytosol and is imported into peroxisomes as a folded protein [96]. Alcohol oxidase from methylotrophic yeasts also binds its cofactor in the cytosol, although in this case the assembly into an active octamer occurs inside peroxisomes [97,98]. Thus it appears likely that peroxisomal FAD-containing oxidases generally acquire their cofactor in the cytosol, and a peroxisomal FAD transporter might not be required. Peroxisomal FAD is reoxidised with the concomitant generation of H_2O_2 which in turn is removed by catalase. This situation contrasts with the import of FAD into mitochondria: the mitochondrial inner membrane has at least two specific FAD transporters, with Flx1p representing a FAD exporter [99], but in this organelle proteins traverse the membrane in an unfolded state which necessitates an intramitochondrial cofactor charging of the apo-flavoenzymes.

NADH is produced in the course of β -oxidation by the dehydrogenase reaction of the multifunctional enzyme and must be regenerated via one or more redox shuttles analogous to those in mitochondria. In mammals, pyruvate or dihydroxyacetone phosphate are likely to serve as the electron acceptors, through the action of lactate dehydrogenase [100,101] and glycerol-3-phosphate dehydrogenase [102], respectively. In contrast, in yeast [10] and plants (Pracharoenwattana and Smith, personal communication), the electrons are transferred to oxaloacetate, which is converted to malate by peroxisomal malate dehydrogenase. This could operate in the context of two different shuttles: a malate-oxaloacetate shuttle, or a malate-aspartate shuttle in which reducing equivalents are transferred from peroxisomes to mitochondria [10,103]. The former is a simpler scenario in which malate is exported to the cytoplasm, where it is re-oxidised by cytosolic malate dehydrogenase to oxaloacetate, which re-enters the peroxisome (Fig. 2A). NADH produced by cytosolic malate dehydrogenase is then re-oxidised by mitochondrial NADH dehydrogenases. The alternative malate-aspartate shuttle was originally proposed due to the fact that the inner mitochondrial membrane is relatively impermeable to oxaloacetate. In this case, oxaloacetate is not imported into the peroxisome, but rather is generated from aspartate and 2-oxoglutarate with the concomitant formation of glutamate in a transamination reaction catalysed by aspartate amino transferase [103,104]. This requires exchange of malate/2-oxoglutarate and glutamate/aspartate across the peroxisomal and mitochondrial membranes (Fig. 2B). However, this scheme has been contested by biochemical studies with leaf peroxisomes [105] and disruption of the *AAT2* gene of yeast does not




Fig. 2. Transport of reducing equivalents. The dehydrogenase reaction of the multifunctional protein of β -oxidation requires NAD^+ . Since the peroxisomal membrane is impermeable to nucleotides and nucleotide-containing cofactors, two redox shuttles involved in regeneration of peroxisomal NAD^+ have been proposed for plants and yeast. Both systems require exchange of organic anions across the peroxisomal membrane, which may be executed by specific carriers or porin-like proteins. (A) The malate-oxaloacetate shuttle. Peroxisomal malate dehydrogenase (MDH) converts oxaloacetate (OAA) to malate with the concomitant oxidation of NADH . Malate is exported to the cytosol, where it is re-converted to OAA by the action of cytosolic MDH. NADH produced in this reaction is re-oxidized by mitochondrial NADH dehydrogenases (NDH). Import of OAA into the peroxisome completes the shuttle. Abbreviations: ACX, acyl-CoA oxidase; MFP, multifunctional protein; thio, thiolase. (B) The malate-2-oxoglutarate shuttle. As in the malate-oxaloacetate shuttle, malate produced from OAA by peroxisomal MDH is exported from the peroxisome. Malate is then transported into mitochondria where it is converted to OAA by MDH. OAA then undergoes conversion to aspartate by aspartate amino transferase (AAT), with the concomitant formation of 2-oxoglutarate (2-OG) from glutamate. 2-OG is transferred from mitochondria to peroxisomes where the reverse reaction occurs, catalysed by peroxisomal aspartate amino transferase, to yield OAA. This shuttle also requires the transport of aspartate and glutamate between mitochondria, peroxisomes and cytosol. Redrawn from Elgersma and Tabak [154].

affect growth on oleate, indicating that it is not essential for the regeneration of intraperoxisomal NAD^+ [106].

Regardless of the specific redox shuttle involved in regeneration, in all cases, NAD(H) must move back and forth from the multifunctional enzyme to a dehydrogenase and is

therefore only transiently bound to the enzymes. It might be that NAD^+ is imported in the manner of FAD through cytosolic association and subsequent joint transit with NAD^+ -dependent enzymes. Whether such a mechanism would be sufficient to establish the peroxisomal pool of NAD(H) remains unclear. It appears more likely that additional, low efficiency routes for provision of peroxisomal NAD^+ exist.

Similar arguments account for the supply of peroxisomes with NADP^+ . This cofactor is required for the β -oxidation auxiliary enzyme Δ^2 - Δ^4 -dienoyl-CoA reductase, which is essential for the degradation of unsaturated fatty acids with double bonds at even-numbered positions [107]. NADPH is regenerated inside peroxisomes via an isocitrate/2-oxoglutarate shuttle [108–110], similar to that which links the mitochondrial and cytosolic pools of NADPH in rat liver (Fig. 3). Oxidation of isocitrate is catalysed by the peroxisomal NADP^+ -dependent isocitrate dehydrogenase, Idp3p. This reaction regenerates NADPH and yields 2-oxoglutarate which is exported from the peroxisome and is possibly re-reduced to isocitrate by the cytosolic isocitrate dehydrogenase, Idp2p. The reverse reductive carboxylation of 2-oxoglutarate appears feasible under physiological conditions, since the K_m values of Idp2p are similar for isocitrate and 2-oxoglutarate, albeit V_{max} is about 20 times lower for the latter substrate [111]. The shuttle is thought to be completed by the import of isocitrate into the peroxisome [109,110], but the carriers required for shuttling of isocitrate and 2-oxoglutarate remain elusive. An alternative route for the peroxisomal regeneration of NADPH involves glucose-6-phosphate dehydrogenase [112].

The operation of an isocitrate/2-oxoglutarate shuttle or a glucose-6-phosphate dehydrogenase-dependent mechanism for NADPH regeneration in plant peroxisomes still requires experimental confirmation, although genes encoding a number of the requisite components have been identified at least *in silico* [104]. However, the identity of the dicarboxylate carriers which function in peroxisomal redox shuttles remains obscure. Mitochondria contain several dicarboxylate carriers which might co-target to peroxisomes [86]. Small amounts of specific carriers might have escaped detection in the various proteomic approaches aimed at identifying peroxisomal membrane proteins due to the prevalent problem of mitochondrial contamination [113,114]. Alternatively, peroxisomal porins which have not yet been identified at the molecular level could be responsible for transport of shuttle metabolites (see Antonenkov and Hiltunen, this issue).

2.5. Nudix hydrolases and regulation of cofactor and co substrate concentration

In addition to import and regeneration, export must be considered as a factor affecting the intraperoxisomal pools of NAD(P)H and CoA. Nudix hydrolases comprise a diverse family of enzymes which hydrolyse a diphosphate linkage in a variety of nucleotide-containing compounds and appear to play an important role in the regulation of intraperoxisomal cofactor concentration and possibly removal of damaged cofactors [115]. Peroxisomal representatives identified to date include




Fig. 3. The isocitrate-2-oxoglutarate shuttle. In addition to the chain-shortening β -oxidation system, the oxidation of unsaturated fatty acids requires auxiliary enzymes for the elimination of double bonds, including Δ^2 - Δ^4 -dienoyl-CoA reductase (DECR) and Δ^3 - Δ^2 -enoyl-CoA isomerase which may be provided by multifunctional protein (MFP) or by a monofunctional peroxisomal Δ^3 - Δ^2 -enoyl-CoA isomerase (PECI). The requirement of DECR for NADPH requires a system for the regeneration of peroxisomal NADPH. The isocitrate-2-oxoglutarate shuttle which employs peroxisomal (Idp3p) and cytosolic (Idp2p) isocitrate dehydrogenase has been proposed to occur in yeast and may also occur in plants. The mechanism of isocitrate and 2-oxo-glutarate transport across the peroxisomal membrane is unknown, but may involve specific carriers or porins which have been shown to transport dicarboxylic acids. Abbreviations: ACX, acyl-CoA oxidase; 2-OG, 2-oxoglutarate. Modified from van Roermund et al. [110].

NAD(P)H diphosphatases which exhibit substrate preference for reduced dinucleotides [116,117] and CoA diphosphatases, which handle CoA, CoA esters and oxidised CoA with differing efficiency [118–121]. Importantly from a transport perspective, these enzymes have the net effect of degrading bulky cofactors to smaller molecules which can readily exit the peroxisome.

3. Export of acetyl- and acyl-units

The export of acetyl-CoAs generated by peroxisomal β -oxidation is accomplished by either of two systems, a carnitine shuttle or through the export of citrate. This is because acetyl-CoA is not able to readily traverse the peroxisomal membrane, whereas the compounds produced are. The relative importance of the two modes varies between species. Genetic evidence indicates that both systems operate in yeast [10], whereas in plants only citrate export occurs [122]. Mammals lack the glyoxylate cycle and a peroxisomal citrate synthase does not exist, suggesting that here the carnitine-mediated export is the exclusive mode of acetyl-removal. An alternative route, which predominates in some mammalian tissues, is the export of acetate generated from acetyl-CoA by the action of peroxisomal thioesterases [1].

3.1. The carnitine shuttle

The existence of a carnitine shuttle is supported by the localisation of yeast carnitine acetyl transferase Cat2p, which

is found in both mitochondria and peroxisomes [123]. In peroxisomes, the enzyme is thought to catalyse transfer of the acetyl moiety from CoA to carnitine, whereas in mitochondria it would catalyse the reverse reaction thereby enabling the condensation of acetyl-CoA with oxaloacetate to form citrate. As a consequence, acetyl-carnitine has to traverse both the peroxisomal as well as the inner mitochondrial membrane. Mitochondrial transport is accomplished by the carnitine carrier Crc1p, a member of the mitochondrial carrier family, which transports acetyl-(and propionyl-) carnitine in exchange for carnitine [124,125]. The important role the *CAT2* and *CRC1* genes play in fatty acid utilisation is also indicated by the induction of their expression in the presence of oleate [126]. However, and in contrast to the dually localised Cat2p transferase, the carnitine carrier is apparently absent from the yeast peroxisomal membrane [125]. This is also true for the *Aspergillus nidulans* carnitine carrier AcuH [127,128]. Which peroxisomal protein therefore enables the exit of acetyl-carnitine and the import of carnitine remains elusive.

In mammals, carnitine octanoyl transferase with a substrate preference for MCFAs additionally resides within peroxisomes [1,129,130]. This enzyme is required because peroxisomal β -oxidation usually does not go to completion. Rather, the chain-shortened substrates are transported to mitochondria as carnitine conjugates, where they are reconverted into the CoA esters and further degraded by the resident β -oxidation system. The broader substrate specificity of the mitochondrial carnitine acylcarnitine carrier CAC, which also accepts longer carnitine fatty acyl moieties, reflects the distinct distribution of β -oxidation systems in mammals and yeast [131]. A cluster of three carnitine transporters has been identified in the mouse genome with distinct subcellular localisation and affinities for carnitine. Recently, it has been suggested that *OCTN3* encodes a peroxisomal medium-affinity carnitine transporter [132,133]. Fusing *OCTN3* to green fluorescent protein (GFP) and immunogold labelling using anti-*OCTN3* antibodies revealed that this isoform of the family might indeed be a predominantly peroxisomal protein. The antibodies also reacted with a protein in human fibroblasts that was absent in a *pex19* patient cell line, in which the steady state concentration of several PMPs is below the detection limit due to a specific defect in peroxisome membrane biogenesis [134]. In light of the fact that the *OCTN* transporters belong to the organic cation transporter family that is not related to the MCF, the relevance of a reported cross-reactivity of an antibody against the mitochondrial carnitine carrier (CAC) with a protein from purified rat peroxisomes remains unclear [135].

Intriguingly, a carnitine acyl carrier homologue, BOU, has also been identified in *Arabidopsis* mitochondria. *bou* mutant plants exhibit a striking phenotype in which growth is arrested after germination but lipids are still degraded [136]. However, the function of this transporter in relation to peroxisomal metabolism remains obscure, since higher plants possess neither peroxisomal carnitine transferases nor a mitochondrial β -oxidation system.

3.2. Carnitine-independent export from peroxisomes: the role of thioesterases

In mammals, a carnitine-independent route for exporting acyl units from peroxisomes has been demonstrated [1,137]. Here, CoA esters are hydrolysed within peroxisomes to yield free acids which are transported to mitochondria and re-activated to their respective CoA esters for completion of β -oxidation. Hydrolysis of CoA esters is catalysed by peroxisomal thioesterases which vary in substrate specificity [138].

Interestingly, the *Arabidopsis* genome contains several candidate peroxisomal thioesterases [104,139], one of which (ACH2) has already been characterised [140]. In addition to roles in fatty acid β -oxidation, these may play an important role in the export of diverse, plant-specific metabolites synthesised in the peroxisome. For example, the final biosynthetic steps of the carboxylate-containing plant hormones, jasmonic acid and indole acetic acid, occur in peroxisomes. Following one (IBA to IAA) or three (OPDA to jasmonic acid) complete rounds of peroxisomal β -oxidation, the hormones are thought to leave the

organelle as free acids, particularly in light of the absence of peroxisomal carnitine transferases. This would require a thioesterase-catalysed release from the CoA ester. Additional roles proposed for peroxisomal thioesterases include the regulation of acyl-CoA/free CoA levels and the prevention of CoA sequestration by intermediates which are β -oxidised inefficiently [138].

3.3. Diversion of acetyl-CoA through citrate synthase; transport requirements of the glyoxylate cycle

Citrate synthase catalyses the formation of citrate from oxaloacetate and acetyl-CoA with the liberation of free CoA. Yeast peroxisomal citrate synthase Cit2p also participates in the glyoxylate cycle, but is not essential for growth on nonfermentable carbon sources since the mitochondrial TCA cycle isoforms Cit1p and/or Cit3p can substitute for Cit2p (for more details, see Kunze et al., this issue). Acetyl-CoA generated within peroxisomes during β -oxidation also does not accumulate in a *cit2* deletion because of the redundant carnitine




Fig. 4. Transport requirements of the glyoxylate cycle. The enzymes of the glyoxylate cycle are located in both peroxisomes and cytosol and isocitrate lyase is differentially localised in plants and fungi. This requires the transport of several different metabolites across the peroxisomal membrane but the precise biochemical nature of the transport steps in the glyoxylate cycle is unknown. Note that this scheme requires cytosolic malate dehydrogenase (MDH); peroxisomal MDH does not participate in the glyoxylate cycle, but rather operates in the reverse direction, in the regeneration of reducing equivalents shown in Fig. 2. The export of malate and import of oxaloacetate (OAA) correspond to the malate-OAA shuttle (Fig. 2A). Abbreviations: CSY, citrate synthase; ACO, aconitase; ICL, isocitrate lyase; MLS, malate synthase.

conjugation system described above. However, by deleting both *CIT2* and *CAT2* or *CRC1*, growth on oleate is abolished, because now acetyl-CoA is trapped within peroxisomes.

In *Arabidopsis*, two glyoxysomal citrate synthases *Csy2* and *Csy3* fulfil an analogous function in β -oxidation, with the difference that a carnitine bypass does not operate [122]. The double *csy2*, *csy3* mutant therefore exhibits an extremely severe phenotype: like *cts* and other strong β -oxidation alleles [31,141], *csy2*, *csy3* seeds are dormant and will only germinate when the seed coat is removed and sucrose is supplied. Moreover, rescued seedlings do not develop further when transferred to soil, indicating the importance of citrate export in plant growth and development [122].

Citrate produced by peroxisomal metabolism is exported to the cytosol and ultimately enters the mitochondrial TCA cycle, but also participates in the glyoxylate cycle to provide C4 biosynthetic precursors. The operation of the glyoxylate cycle requires the transport of intermediates across the peroxisomal membrane since enzymes of the glyoxylate cycle are located both inside and outside the peroxisome (Kunze et al., this issue). The differential localisation of enzymes in plants and fungi results in different transport requirements for the glyoxylate cycle: export of citrate and malate and import of oxaloacetate are common to both plants and fungi, but organisms in which isocitrate lyase is cytosolic must import glyoxylate whereas organisms with peroxisomal isocitrate lyase require the import of isocitrate and export of succinate (Fig. 4). The transport requirements of the glyoxylate cycle are considered in detail by Kunze et al. (this issue).

4. Intraperoxisomal pH

The existence of a proton gradient across the peroxisomal membrane is a long-standing matter of debate and relates to the question as to whether peroxisomes are impermeable to small molecules *in vivo*. Arguments in favour of a peroxisomal pore with an exclusion size of about 300 Da are presented by Antonenkov and Hiltunen (this issue). Data supporting the closed-compartment hypothesis emerged from analyses that were concerned with the pH of peroxisomes. Early evidence for a proton gradient across the peroxisomal membrane was obtained from studies with yeast. Upon induction of *Hansenula polymorpha* peroxisomes with methanol, a second phosphate pool became discernible through ^{31}P NMR studies measuring the pH-dependent chemical shifts of phosphate: in addition to the cytosolic phosphate peak, a peroxisomal peak attributable to a lower pH was observed [142]. Furthermore, the weak base DAMP, which accumulates specifically in acidic compartments, was detectable in peroxisomes of intact spheroplasts by immunocytochemistry [143]. Dissipation of intracellular pH gradients by chloroquine resulted in the disappearance of the second phosphate peak and DAMP was no longer located in peroxisomes, consistent with the existence of an endogenous ΔpH across the peroxisomal membrane. At that time, the generation of a proton gradient was explained by a (hypothetical) peroxisomal proton-translocating ATPase [144,145], but the molecular identity of this activity has remained elusive.

More recent *in vivo* measurements of the peroxisomal pH have caused considerable confusion. Determination of the peroxisomal pH in human fibroblasts with a membrane-permeable peptide harbouring a C-terminal peroxisomal targeting signal type 1 (PTS1) and labelled with the pH-sensitive fluorescent moiety carboxyseminalphtho fluorescein (SNAFL-2) confirmed the existence of a pH gradient, but apparently with a reversed orientation, i.e. basic inside, with a peroxisomal pH of 8.2 [146]. The gradient could be dissipated by the addition of uncoupler and was apparently absent in a cell line harbouring a mutated *PEX7*, which exhibits a specific import defect of proteins with a PTS2. The approach used relies on the pH-dependent emission ratio of the fluorophore at two different wavelengths and is principally well suited for such measurements. A limitation of the method might arise from the low intracellular concentration of peptides dispersed throughout the cytosol, making cytosolic pH measurements troublesome. Its general applicability is further restricted by the possible extrusion of these peptides via multiple drug resistance (MDR) transporters [147] and the uptake of such peptides via endocytosis in yeast (our unpublished observation).

Variants of GFP allow a similar detection of pH-dependent emission intensities with the advantage of intracellular protein expression. Expression of one of these variants, the so-called ratiometric pHluorin [148] as an SKL-tagged variant in Chinese hamster ovary (CHO) and human fibroblasts revealed that the peroxisomal pH adapts to that of the cytosol [149]. As a consequence, the compartment was classified as being proton-permeable, not taking into account that changing a gradient on one side evokes an adjustment on the other side of the membrane. In apparent contrast, by expressing the same protein in yeast, a pH value more acidic than that of the cytosol was determined for peroxisomes [88]. Similar emission ratios of ratiometric pHluorin were observed for cytosolic and peroxisomal variants in both systems, but attempts to calibrate the system from outside the cell by Jankowski et al. [149] led to a different interpretation of the results.

Yellow fluorescent protein (YFP) and a point-mutated version with an altered pKa value were also used to determine the pH of yeast peroxisomes. Here, the absolute fluorescence intensity (rather than a ratio of intensities) was correlated with the pH, which possibly makes the system more vulnerable to differences in expression and distribution of the fluorescent protein. The outcome of this study was an alkaline peroxisomal pH that was even higher in several mutants affected in β -oxidation, whereas in *ant1* Δ cells, a pH gradient was not observed. A hypothesis was put forward that the imposed pH gradient serves as a driving force for the passive transport of free MCFAs, which would deprotonate more readily inside peroxisomes in a basic milieu [89]. However, this idea is hard to reconcile with the observed depletion of the gradient in the absence of Ant1p, which increases the luminal proton concentration, unless an intraperoxisomal ATP-consuming reaction or transport process overcompensates this effect. Support for a basic pH also enters from studies with *Penicillium chrysogenum* suggesting a slightly alkaline pH for the microbody lumen in this fungus [150]. Finally, the basic pI of

several peroxisomal matrix enzymes has been invoked to support an argument for an alkaline peroxisomal pH [146].

The jury is out there to track down possible caveats in the various methods applied, which should lead to the desirable clarification of this issue. An important factor to be considered in future studies is the influence of growth conditions, developmental state and cell type on the peroxisomal pH. For instance, methanol utilisation in *H. polymorpha* requires an acidic peroxisomal pH, since the substrate specificity of dihydroxyacetone synthase requires formaldehyde in its protonated form [151]. What seems clear is that in yeast, Ant1p contributes to the formation of a pH gradient as a direct consequence of its proton-compensated transport mode [88]. Although the existence of an H⁺ATPase is unlikely, additional factors might contribute to the maintenance of this gradient. To date it can only be speculated whether a peroxisomal proton gradient is needed to drive other transport processes, though physicochemical considerations predict that the existence of a pH gradient between the cytosol and peroxisomal lumen is likely to be a key component in determining the extent of passive transport of free acids across the peroxisomal membrane.

5. Conclusions and perspectives

Although considerable progress in understanding of peroxisomal transport processes has been made in recent years, many questions remain to be resolved. Several novel peroxisomal transporters have been reported recently, including a Ca²⁺ATPase in *Drosophila* [152] and a peroxisomal protein from brome grass (*Bromus inermis*) which is similar to amino acid-selective channel proteins from the chloroplast outer membrane [153]. Nevertheless, knowledge of peroxisomal metabolism predicts the existence of further metabolite transporters involved in organism-specific peroxisomal processes, in addition to those mentioned elsewhere in this review. Identification of such transporters may come from several approaches, including proteomic analysis of peroxisomal membrane proteins, forward genetics approaches and characterisation of candidate genes using knock-down systems. A further, yet exciting, challenge is the rigorous characterisation of known transporters which will require the development of transport assays with isolated organelles and reconstituted, purified proteins. The past few decades have been an exciting era for peroxisome research and an improved understanding of peroxisomal transporters will undoubtedly enhance and extend our appreciation of the functions of this multipurpose organelle.

Acknowledgements

Work on peroxisomal ABC transporters in FLT's laboratory is supported by Biology and Biotechnology Research Council (BBSRC) grant P19769. Rothamsted Research receives grant-aided support from the BBSRC of the UK. The work of HR's laboratory is supported by the Deutsche Forschungsgemeinschaft grants SFB480 and RO2494/1.

References

- [1] R.J. Wanders, H.R. Waterham, Biochemistry of mammalian peroxisomes revisited, *Annu. Rev. Biochem.* 75 (2006) 295–332.
- [2] M. Hayashi, M. Nishimura, Entering a new era of research on plant peroxisomes, *Curr. Opin. Plant Biol.* 6 (2003) 577–582.
- [3] F.L. Theodoulou, M. Holdsworth, A. Baker, Peroxisomal ABC transporters, *FEBS Lett.* 580 (2006) 1139–1155.
- [4] V.I. Titorenko, R.A. Rachubinski, The peroxisome: orchestrating important developmental decisions from inside the cell, *J. Cell Biol.* 164 (2004) 641–645.
- [5] T. Kamada, K. Nito, H. Hayashi, S. Mano, M. Hayashi, M. Nishimura, Functional differentiation of peroxisomes revealed by expression profiles of peroxisomal genes in *Arabidopsis thaliana*, *Plant Cell Physiol.* 44 (2003) 1275–1289.
- [6] P.A. Michels, V. Hannaert, F. Bringaud, Metabolic aspects of glycosomes in trypanosomatidae—new data and views, *Parasitol. Today* 16 (2000) 482–489.
- [7] M. Parsons, Glycosomes: parasites and the divergence of peroxisomal purpose, *Mol. Microbiol.* 53 (2004) 717–724.
- [8] S. Reumann, The structural properties of plant peroxisomes and their metabolic significance, *Biol. Chem.* 381 (2000) 639–648.
- [9] V.D. Antonenkov, R.T. Sormunen, J.K. Hiltunen, The rat liver peroxisomal membrane forms a permeability barrier for cofactors but not for small metabolites in vitro, *J. Cell Sci.* 117 (2004) 5633–5642.
- [10] C.W. van Roermund, Y. Elgersma, N. Singh, R.J. Wanders, H.F. Tabak, The membrane of peroxisomes in *Saccharomyces cerevisiae* is impermeable to NAD(H) and acetyl-CoA under in vivo conditions, *EMBO J.* 14 (1995) 3480–3486.
- [11] A. Baker, I.A. Graham, M. Holdsworth, S.M. Smith, F.L. Theodoulou, Chewing the fat: β -oxidation in signalling and development, *Trends Plant Sci.* 11 (2006) 124–132.
- [12] C.F. Higgins, ABC transporters: from microorganisms to man, *Annu. Rev. Cell Biol.* 8 (1992) 67–113.
- [13] C. Anjard, W.F. Loomis, Evolutionary analyses of ABC transporters of *Dictyostelium discoideum*, *Eukaryot. Cell* 1 (2002) 643–652.
- [14] M. Dean, A. Rzhetsky, R. Allikmets, The human ATP-binding cassette (ABC) transporter superfamily, *Genome Res.* 11 (2001) 1156–1166.
- [15] S. Footitt, S.P. Slocombe, V. Larner, S. Kurup, Y. Wu, T. Larson, I. Graham, A. Baker, M. Holdsworth, Control of germination and lipid mobilization by COMATOSE, the *Arabidopsis* homologue of human ALDP, *EMBO J.* 21 (2002) 2912–2922.
- [16] O. Garcia, P. Bouige, C. Forestier, E. Dassa, Inventory and comparative analysis of rice and *Arabidopsis* ATP-binding cassette (ABC) systems, *J. Mol. Biol.* 343 (2004) 249–265.
- [17] H. Jungwirth, K. Kuchler, Yeast ABC transporters—A tale of sex, stress, drugs and aging, *FEBS Lett.* 580 (2006) 1131–1138.
- [18] O.I. Petriv, D.B. Pilgrim, R.A. Rachubinski, V.I. Titorenko, RNA interference of peroxisome-related genes in *C. elegans*: a new model for human peroxisomal disorders, *Physiol. Genomics* 10 (2002) 79–91.
- [19] C.W. Roth, I. Holm, M. Graille, P. Dehoux, A. Rzhetsky, P. Wincker, J. Weissenbach, P.T. Brey, Identification of the *Anopheles gambiae* ATP-binding cassette transporter superfamily genes, *Mol. Cells* 15 (2003) 150–158.
- [20] L.M. Schriml, M. Dean, Identification of 18 mouse ABC genes and characterization of the ABC superfamily in *Mus musculus*, *Genomics* 64 (2000) 24–31.
- [21] J.A. Sheps, S. Ralph, Z. Zhao, D.L. Baillie, V. Ling, The ABC transporter gene family of *Caenorhabditis elegans* has implications for the evolutionary dynamics of multidrug resistance in eukaryotes, *Genome Biol.* 5 (2004) R15.
- [22] C. Yernaux, M. Fransen, C. Brees, S. Lorenzen, P.A. Michels, *Trypanosoma brucei* glycosomal ABC transporters: identification and membrane targeting, *Mol. Membr. Biol.* 23 (2006) 157–172.
- [23] E.H. Hetteema, C.W. van Roermund, B. Distel, M. van den Berg, C. Vilela, C. Rodrigues-Pousada, R.J. Wanders, H.F. Tabak, The ABC transporter proteins Pat1 and Pat2 are required for import of long-chain

- fatty acids into peroxisomes of *Saccharomyces cerevisiae*, *EMBO J.* 15 (1996) 3813–3822.
- [24] N. Shani, P.A. Watkins, D. Valle, *PXA1*, a possible *Saccharomyces cerevisiae* ortholog of the human adrenoleukodystrophy gene, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 6012–6016.
- [25] N. Shani, D. Valle, A *Saccharomyces cerevisiae* homolog of the human adrenoleukodystrophy transporter is a heterodimer of two half ATP-binding cassette transporters, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 11901–11906.
- [26] E.E. Swartzman, M.N. Viswanathan, J. Thorer, The *PAL1* gene product is a peroxisomal ATP-binding cassette transporter in the yeast *Saccharomyces cerevisiae*, *J. Cell Biol.* 132 (1996) 549–563.
- [27] N. Verleur, E.H. Hettema, C.W. van Roermund, H.F. Tabak, R.J. Wanders, Transport of activated fatty acids by the peroxisomal ATP-binding-cassette transporter Pxa2 in a semi-intact yeast cell system, *Eur. J. Biochem.* 249 (1997) 657–661.
- [28] E.H. Hettema, H.F. Tabak, Transport of fatty acids and metabolites across the peroxisomal membrane, *Biochim. Biophys. Acta* 1486 (2000) 18–27.
- [29] B.K. Zolman, I.D. Silva, B. Bartel, The *Arabidopsis* pxa1 mutant is defective in an ATP-binding cassette transporter-like protein required for peroxisomal fatty acid β -oxidation, *Plant Physiol.* 127 (2001) 1266–1278.
- [30] M. Hayashi, K. Nito, R. Takei-Hoshi, M. Yagi, M. Kondo, A. Suenaga, T. Yamaya, M. Nishimura, Ped3p is a peroxisomal ATP-binding cassette transporter that might supply substrates for fatty acid β -oxidation, *Plant Cell Physiol.* 43 (2002) 1–11.
- [31] S. Footitt, J. Marquez, H. Schmuths, A. Baker, F.L. Theodoulou, M. Holdsworth, Analysis of the role of *COMATOSE* and peroxisomal β -oxidation in the determination of germination potential in *Arabidopsis*, *J. Exp. Bot.* 57 (2006) 2805–2814.
- [32] M. Fulda, J. Schnurr, A. Abbadi, E. Heinz, J. Browse, Peroxisomal Acyl-CoA synthetase activity is essential for seedling development in *Arabidopsis thaliana*, *Plant Cell* 16 (2004) 394–405.
- [33] M. Fulda, J. Shockey, M. Werber, F.P. Wolter, E. Heinz, Two long-chain acyl-CoA synthetases from *Arabidopsis thaliana* involved in peroxisomal fatty acid β -oxidation, *Plant J.* 32 (2002) 93–103.
- [34] V. Germain, E.L. Rylott, T.R. Larson, S.M. Sherson, N. Bechtold, J.P. Carde, J.H. Bryce, I.A. Graham, S.M. Smith, Requirement for 3-ketoacyl-CoA thiolase-2 in peroxisome development, fatty acid β -oxidation and breakdown of triacylglycerol in lipid bodies of *Arabidopsis* seedlings, *Plant J.* 28 (2001) 1–12.
- [35] E.L. Rylott, P.J. Eastmond, A.D. Gilday, S.P. Slocombe, T.R. Larson, A. Baker, I.A. Graham, The *Arabidopsis thaliana* multifunctional protein gene (MFP2) of peroxisomal β -oxidation is essential for seedling establishment, *Plant J.* 45 (2006) 930–941.
- [36] I.A. Graham, Y. Li, T.R. Larson, Acyl-CoA measurements in plants suggest a role in regulating various cellular processes, *Biochem. Soc. Trans.* 30 (2002) 1095–1099.
- [37] F.L. Theodoulou, K. Job, S.P. Slocombe, S. Footitt, M. Holdsworth, A. Baker, T.R. Larson, I.A. Graham, Jasmonic acid levels are reduced in *COMATOSE* ATP-binding cassette transporter mutants. Implications for transport of jasmonate precursors into peroxisomes, *Plant Physiol.* 137 (2005) 835–840.
- [38] A. Holzinger, S. Kammerer, A.A. Roscher, Primary structure of human PMP69, a putative peroxisomal ABC-transporter, *Biochem. Biophys. Res. Commun.* 237 (1997) 152–157.
- [39] A. Holzinger, S. Kammerer, J. Berger, A.A. Roscher, cDNA cloning and mRNA expression of the human adrenoleukodystrophy related protein (ALDRP), a peroxisomal ABC transporter, *Biochem. Biophys. Res. Commun.* 239 (1997) 261–264.
- [40] K. Kamijo, S. Taketani, S. Yokota, T. Osumi, T. Hashimoto, The 70-kDa peroxisomal membrane protein is a member of the Mdr (P-glycoprotein)-related ATP-binding protein superfamily, *J. Biol. Chem.* 265 (1990) 4534–4540.
- [41] G. Lombard-Platet, S. Savary, C.O. Sarde, J.L. Mandel, G. Chimini, A close relative of the adrenoleukodystrophy (ALD) gene codes for a peroxisomal protein with a specific expression pattern, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 1265–1269.
- [42] J. Mosser, A.M. Douar, C.O. Sarde, P. Kioschis, R. Feil, H. Moser, A.M. Poustka, J.L. Mandel, P. Aubourg, Putative X-linked adrenoleukodystrophy gene shares unexpected homology with ABC transporters, *Nature* 361 (1993) 726–730.
- [43] N. Shani, G. Jimenez-Sanchez, G. Steel, M. Dean, D. Valle, Identification of a fourth half ABC transporter in the human peroxisomal membrane, *Hum. Mol. Genet.* 6 (1997) 1925–1931.
- [44] M. Contreras, J. Mosser, J.L. Mandel, P. Aubourg, I. Singh, The protein coded by the X-adrenoleukodystrophy gene is a peroxisomal integral membrane protein, *FEBS Lett.* 344 (1994) 211–215.
- [45] J. Mosser, Y. Lutz, M.E. Stoeckel, C.O. Sarde, C. Kretz, A.M. Douar, J. Lopez, P. Aubourg, J.L. Mandel, The gene responsible for adrenoleukodystrophy encodes a peroxisomal membrane protein, *Hum. Mol. Genet.* 3 (1994) 265–271.
- [46] P.A. Watkins, S.J. Gould, M.A. Smith, L.T. Braiterman, H.M. Wei, F. Kok, A.B. Moser, H.W. Moser, K.D. Smith, Altered expression of ALDP in X-linked adrenoleukodystrophy, *Am. J. Hum. Genet.* 57 (1995) 292–301.
- [47] C.J. Gloeckner, P.U. Mayerhofer, P. Landgraf, A.C. Muntau, A. Holzinger, J.K. Gerber, S. Kammerer, J. Adamski, A.A. Roscher, Human adrenoleukodystrophy protein and related peroxisomal ABC transporters interact with the peroxisomal assembly protein PEX19p, *Biochem. Biophys. Res. Commun.* 271 (2000) 144–150.
- [48] Y. Kashiwayama, K. Asahina, H. Shibata, M. Morita, A.C. Muntau, A.A. Roscher, R.J. Wanders, N. Shimozawa, M. Sakaguchi, H. Kato, T. Imanaka, Role of Pex19p in the targeting of PMP70 to peroxisome, *Biochim. Biophys. Acta* 1746 (2005) 116–128.
- [49] A. Halbach, S. Lorenzen, C. Landgraf, R. Volkmer-Engert, R. Erdmann, H. Rottensteiner, Function of the PEX19-binding site of human adrenoleukodystrophy protein as targeting motif in man and yeast: PMP targeting is evolutionarily conserved, *J. Biol. Chem.* 280 (2005) 21176–21182.
- [50] J.M. Jones, J.C. Morrell, S.J. Gould, PEX19 is a predominantly cytosolic chaperone and import receptor for class I peroxisomal membrane proteins, *J. Cell Biol.* 164 (2004) 57–67.
- [51] T. Imanaka, K. Aihara, T. Takano, A. Yamashita, R. Sato, Y. Suzuki, S. Yokota, T. Osumi, Characterization of the 70-kDa peroxisomal membrane protein, an ATP binding cassette transporter, *J. Biol. Chem.* 274 (1999) 11968–11976.
- [52] R.J. Wanders, Peroxisomes, lipid metabolism, and peroxisomal disorders, *Mol. Genet. Metab.* 83 (2004) 16–27.
- [53] R.J. Wanders, P. Vreken, S. Ferdinandusse, G.A. Jansen, H.R. Waterham, C.W. van Roermund, E.G. Van Grunsven, Peroxisomal fatty acid α - and β -oxidation in humans: enzymology, peroxisomal metabolite transporters and peroxisomal diseases, *Biochem. Soc. Trans.* 29 (2001) 250–267.
- [54] I. Silva-Zolezzi, K. Hebron, S. Mihalik, G. Thomas, D. Valle, G. Jimenez-Sanchez, 53rd Annual Meeting of The American Society of Human Genetics, Los Angeles, CA, 2003.
- [55] M. Hashmi, W. Stanley, I. Singh, Lignoceryl-CoA-SH ligase: enzyme defect in fatty acid β -oxidation system in X-linked childhood adrenoleukodystrophy, *FEBS Lett.* 196 (1986) 247–250.
- [56] O. Lazo, M. Contreras, M. Hashmi, W. Stanley, C. Irazu, I. Singh, Peroxisomal lignoceryl-CoA ligase deficiency in childhood adrenoleukodystrophy and adrenomyeloneuropathy, *Proc. Natl. Acad. Sci. U. S. A.* 85 (1988) 7647–7651.
- [57] R.J. Wanders, C.W. van Roermund, M.J. van Wijland, R.B. Schutgens, J. Heikooop, H. van den Bosch, A.W. Schram, J.M. Tager, Peroxisomal fatty acid β -oxidation in relation to the accumulation of very long-chain fatty acids in cultured skin fibroblasts from patients with Zellweger syndrome and other peroxisomal disorders, *J. Clin. Invest.* 80 (1987) 1778–1783.
- [58] L.T. Braiterman, S. Zheng, P.A. Watkins, M.T. Geraghty, G. Johnson, M.C. McGuinness, A.B. Moser, K.D. Smith, Suppression of peroxisomal membrane protein defects by peroxisomal ATP binding cassette (ABC) proteins, *Hum. Mol. Genet.* 7 (1998) 239–247.
- [59] N. Shinnoh, T. Yamada, T. Yoshimura, H. Furuya, Y. Yoshida, Y. Suzuki, N. Shimozawa, T. Orii, T. Kobayashi, Adrenoleukodystrophy: the restoration of peroxisomal β -oxidation by transfection of normal cDNA, *Biochem. Biophys. Res. Commun.* 210 (1995) 830–836.

- [60] N. Cartier, J. Lopez, P. Moullier, F. Rocchiccioli, M.O. Rolland, P. Jorge, J. Mosser, J.L. Mandel, P.F. Bougneres, O. Danos, et al., Retroviral-mediated gene transfer corrects very-long-chain fatty acid metabolism in adrenoleukodystrophy fibroblasts, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 1674–1678.
- [61] V. Feigenbaum, G. Lombard-Platet, S. Guidoux, C.O. Sarde, J.L. Mandel, P. Aubourg, Mutational and protein analysis of patients and heterozygous women with X-linked adrenoleukodystrophy, *Am. J. Hum. Genet.* 58 (1996) 1135–1144.
- [62] A. Dodd, S.A. Rowland, S.L. Hawkes, M.A. Kennedy, D.R. Love, Mutations in the adrenoleukodystrophy gene, *Hum. Mutat.* 9 (1997) 500–511.
- [63] K.D. Smith, S. Kemp, L.T. Braiterman, J.F. Lu, H.M. Wei, M. Geraghty, G. Stetten, J.S. Bergin, J. Pevsner, P.A. Watkins, X-linked adrenoleukodystrophy: genes, mutations, and phenotypes, *Neurochem. Res.* 24 (1999) 521–535.
- [64] S. Kemp, A. Pujol, H.R. Waterham, B.M. van Geel, C.D. Boehm, G.V. Raymond, G.R. Cutting, R.J. Wanders, H.W. Moser, ABCD1 mutations and the X-linked adrenoleukodystrophy mutation database: role in diagnosis and clinical correlations, *Hum. Mutat.* 18 (2001) 499–515.
- [65] T. Kobayashi, N. Shinnoh, A. Kondo, T. Yamada, Adrenoleukodystrophy protein-deficient mice represent abnormality of very long chain fatty acid metabolism, *Biochem. Biophys. Res. Commun.* 232 (1997) 631–636.
- [66] S. Forss-Petter, H. Werner, J. Berger, H. Lassmann, B. Molzer, M.H. Schwab, H. Bernheimer, F. Zimmermann, K.A. Nave, Targeted inactivation of the X-linked adrenoleukodystrophy gene in mice, *J. Neurosci. Res.* 50 (1997) 829–843.
- [67] J.F. Lu, A.M. Lawler, P.A. Watkins, J.M. Powers, A.B. Moser, H.W. Moser, K.D. Smith, A mouse model for X-linked adrenoleukodystrophy, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 9366–9371.
- [68] M.C. McGuinness, J.F. Lu, H.P. Zhang, G.X. Dong, A.K. Heinzer, P.A. Watkins, J. Powers, K.D. Smith, Role of ALDP (ABCD1) and mitochondria in X-linked adrenoleukodystrophy, *Mol. Cell. Biol.* 23 (2003) 744–753.
- [69] I. Oezen, W. Rossmannith, S. Forss-Petter, S. Kemp, T. Voigtlander, K. Moser-Thier, R.J. Wanders, R.E. Bittner, J. Berger, Accumulation of very long-chain fatty acids does not affect mitochondrial function in adrenoleukodystrophy protein deficiency, *Hum. Mol. Genet.* 14 (2005) 1127–1137.
- [70] S. Kemp, F. Valianpour, S. Denis, R. Ofman, R.J. Sanders, P. Mooyer, P.G. Barth, R.J. Wanders, Elongation of very long-chain fatty acids is enhanced in X-linked adrenoleukodystrophy, *Mol. Genet. Metab.* 84 (2005) 144–151.
- [71] C.P. Guimaraes, C. Sa-Miranda, J.E. Azevedo, Probing substrate-induced conformational alterations in adrenoleukodystrophy protein by proteolysis, *J. Hum. Genet.* 50 (2005) 99–105.
- [72] E. Flavigny, A. Sanhaj, P. Aubourg, N. Cartier, Retroviral-mediated adrenoleukodystrophy-related gene transfer corrects very long chain fatty acid metabolism in adrenoleukodystrophy fibroblasts: implications for therapy, *FEBS Lett.* 448 (1999) 261–264.
- [73] S. Kemp, H.M. Wei, J.F. Lu, L.T. Braiterman, M.C. McGuinness, A.B. Moser, P.A. Watkins, K.D. Smith, Gene redundancy and pharmacological gene therapy: implications for X-linked adrenoleukodystrophy, *Nat. Med.* 4 (1998) 1261–1268.
- [74] A. Netik, S. Forss-Petter, A. Holzinger, B. Molzer, G. Unterrainer, J. Berger, Adrenoleukodystrophy-related protein can compensate functionally for adrenoleukodystrophy protein deficiency (X-ALD): implications for therapy, *Hum. Mol. Genet.* 8 (1999) 907–913.
- [75] A. Pujol, I. Ferrer, C. Camps, E. Metzger, C. Hindelang, N. Callizot, M. Ruiz, T. Pampols, M. Giros, J.L. Mandel, Functional overlap between ABCD1 (ALD) and ABCD2 (ALDR) transporters: a therapeutic target for X-adrenoleukodystrophy, *Hum. Mol. Genet.* 13 (2004) 2997–3006.
- [76] D. Valle, J. Gärtner, Human genetics. Penetrating the peroxisome, *Nature* 361 (1993) 682–683.
- [77] G.D. Ewart, D. Cannell, G.B. Cox, A.J. Howells, Mutational analysis of the traffic ATPase (ABC) transporters involved in uptake of eye pigment precursors in *Drosophila melanogaster*. Implications for structure–function relationships, *J. Biol. Chem.* 269 (1994) 10370–10377.
- [78] A.R. Tanaka, K. Tanabe, M. Morita, M. Kurisu, Y. Kasiwayama, M. Matsuo, N. Kioka, T. Amachi, T. Imanaka, K. Ueda, ATP binding/hydrolysis by and phosphorylation of peroxisomal ATP-binding cassette proteins PMP70 (ABCD3) and adrenoleukodystrophy protein (ABCD1), *J. Biol. Chem.* 277 (2002) 40142–40147.
- [79] L.X. Liu, K. Janvier, V. Berteaux-Lecellier, N. Cartier, R. Benarous, P. Aubourg, Homo- and heterodimerization of peroxisomal ATP-binding cassette half-transporters, *J. Biol. Chem.* 274 (1999) 32738–32743.
- [80] N. Troffer-Charlier, N. Doerflinger, E. Metzger, F. Fouquet, J.L. Mandel, P. Aubourg, Mirror expression of adrenoleukodystrophy and adrenoleukodystrophy related genes in mouse tissues and human cell lines, *Eur. J. Cell Biol.* 75 (1998) 254–264.
- [81] I. Ferrer, J.P. Kapfhammer, C. Hindelang, S. Kemp, N. Troffer-Charlier, V. Broccoli, N. Callyzot, P. Mooyer, J. Selhorst, P. Vreken, R.J. Wanders, J.L. Mandel, A. Pujol, Inactivation of the peroxisomal ABCD2 transporter in the mouse leads to late-onset ataxia involving mitochondria, Golgi and endoplasmic reticulum damage, *Hum. Mol. Genet.* 14 (2005) 3565–3577.
- [82] S. Albet, C. Causeret, M. Bentejac, J.L. Mandel, P. Aubourg, B. Maurice, Fenofibrate differently alters expression of genes encoding ATP-binding transporter proteins of the peroxisomal membrane, *FEBS Lett.* 405 (1997) 394–397.
- [83] C.P. Guimaraes, P. Domingues, P. Aubourg, F. Fouquet, A. Pujol, G. Jimenez-Sanchez, C. Sa-Miranda, J.E. Azevedo, Mouse liver PMP70 and ALDP: homomeric interactions prevail in vivo, *Biochim. Biophys. Acta* 1689 (2004) 235–243.
- [84] L. Palmieri, H. Rottensteiner, W. Girzalsky, P. Scarcia, F. Palmieri, R. Erdmann, Identification and functional reconstitution of the yeast peroxisomal adenine nucleotide transporter, *EMBO J.* 20 (2001) 5049–5059.
- [85] C.W. van Roermund, R. Drissen, M. van Den Berg, L. Ijlst, E.H. Hettema, H.F. Tabak, H.R. Waterham, R.J. Wanders, Identification of a peroxisomal ATP carrier required for medium-chain fatty acid β -oxidation and normal peroxisome proliferation in *Saccharomyces cerevisiae*, *Mol. Cell. Biol.* 21 (2001) 4321–4329.
- [86] F. Palmieri, The mitochondrial transporter family (SLC25): physiological and pathological implications, *Pflügers Arch.* 447 (2004) 689–709.
- [87] M. Klingenberg, The ADP, ATP shuttle of the mitochondrion, *Trends Biochem. Sci.* 4 (1979) 249–252.
- [88] F.M. Lasorsa, P. Scarcia, R. Erdmann, F. Palmieri, H. Rottensteiner, L. Palmieri, The yeast peroxisomal adenine nucleotide transporter: characterization of two transport modes and involvement in Δ pH formation across peroxisomal membranes, *Biochem. J.* 381 (2004) 581–585.
- [89] C.W. van Roermund, M. de Jong, I.J. L., J. van Marle, T.B. Dansen, R.J. Wanders, H.R. Waterham, The peroxisomal lumen in *Saccharomyces cerevisiae* is alkaline, *J. Cell Sci.* 117 (2004) 4231–4237.
- [90] T. Nakagawa, T. Imanaka, M. Morita, K. Ishiguro, H. Yurimoto, A. Yamashita, N. Kato, Y. Sakai, Peroxisomal membrane protein Pmp47 is essential in the metabolism of middle-chain fatty acid in yeast peroxisomes and is associated with peroxisome proliferation, *J. Biol. Chem.* 275 (2000) 3455–3461.
- [91] T. Wylin, M. Baes, C. Brees, G.P. Mannaerts, M. Fransen, P.P. Van Veldhoven, Identification and characterization of human PMP34, a protein closely related to the peroxisomal integral membrane protein PMP47 of *Candida boidinii*, *Eur. J. Biochem.* 258 (1998) 332–338.
- [92] W.F. Visser, C.W. van Roermund, H.R. Waterham, R.J. Wanders, Identification of human PMP34 as a peroxisomal ATP transporter, *Biochem. Biophys. Res. Commun.* 299 (2002) 494–497.
- [93] Y. Fukao, Y. Hayashi, S. Mano, M. Hayashi, M. Nishimura, Developmental analysis of a putative ATP/ADP carrier protein localized on glyoxysomal membranes during the peroxisome transition in pumpkin cotyledons, *Plant Cell Physiol.* 42 (2001) 835–841.
- [94] C. Prohl, W. Pelzer, K. Diekert, H. Kmita, T. Bedekovics, G. Kispal, R. Lill, The yeast mitochondrial carrier Leu5p and its human homologue Graves' disease protein are required for accumulation of coenzyme A in the matrix, *Mol. Cell. Biol.* 21 (2001) 1089–1097.
- [95] W.F. Visser, C.W. Van Roermund, L. Ijlst, K.J. Hellingwerf, R.J. Wanders, H.R. Waterham, Demonstration and characterization of

- phosphate transport in mammalian peroxisomes, *Biochem. J.* 389 (2005) 717–722.
- [96] V.I. Titorenko, J.M. Nicaud, H. Wang, H. Chan, R.A. Rachubinski, Acyl-CoA oxidase is imported as a heteropentameric, cofactor-containing complex into peroxisomes of *Yarrowia lipolytica*, *J. Cell Biol.* 156 (2002) 481–494.
- [97] P. Ozimek, M. Veenhuis, I.J. van der Klei, Alcohol oxidase: a complex peroxisomal, oligomeric flavoprotein, *FEMS Yeast Res.* 5 (2005) 975–983.
- [98] M.Q. Stewart, R.D. Esposito, J. Gowani, J.M. Goodman, Alcohol oxidase and dihydroxyacetone synthase, the abundant peroxisomal proteins of methylotrophic yeasts, assemble in different cellular compartments, *J. Cell Sci.* 114 (2001) 2863–2868.
- [99] V. Bafunno, T.A. Giancaspero, C. Brizio, D. Bufano, S. Passarella, E. Boles, M. Barile, Riboflavin uptake and FAD synthesis in *Saccharomyces cerevisiae* mitochondria: involvement of the Flx1p carrier in FAD export, *J. Biol. Chem.* 279 (2004) 95–102.
- [100] G.B. McClelland, S. Khanna, G.F. Gonzalez, C.E. Butz, G.A. Brooks, Peroxisomal membrane monocarboxylate transporters: evidence for a redox shuttle system? *Biochem. Biophys. Res. Commun.* 304 (2003) 130–135.
- [101] E. Baumgart, H.D. Fahimi, A. Stich, A. Völkl, L-lactate dehydrogenase A4- and A3B isoforms are bona fide peroxisomal enzymes in rat liver. Evidence for involvement in intraperoxisomal NADH reoxidation, *J. Biol. Chem.* 271 (1996) 3846–3855.
- [102] R. Gee, E. McGroarty, B. Hsieh, D.M. Wied, N.E. Tolbert, Glycerol phosphate dehydrogenase in mammalian peroxisomes, *Arch. Biochem. Biophys.* 161 (1974) 187–193.
- [103] I.J. Mettler, H. Beevers, Oxidation of NADH in glyoxysomes by a malate-aspartate shuttle, *Plant Physiol.* 66 (1980) 555–560.
- [104] S. Reumann, C. Ma, S. Lemke, L. Babujee, AraPeroX. A database of putative *Arabidopsis* proteins from plant peroxisomes, *Plant Physiol.* 136 (2004) 2587–2608.
- [105] M.R. Schmitt, G.E. Edwards, Provisions of reductant for the hydroxypyruvate to glycerate conversion in leaf peroxisomes: a critical evaluation of the proposed malate/aspartate shuttle, *Plant Physiol.* 72 (1983) 728–734.
- [106] N. Verleur, Y. Elgersma, C.W. Van Roermund, H.F. Tabak, R.J. Wanders, Cytosolic aspartate aminotransferase encoded by the *AAT2* gene is targeted to the peroxisomes in oleate-grown *Saccharomyces cerevisiae*, *Eur. J. Biochem.* 247 (1997) 972–980.
- [107] J.K. Hiltunen, A.M. Mursula, H. Rottensteiner, R.K. Wierenga, A.J. Kastaniotis, A. Gurvitz, The biochemistry of peroxisomal β -oxidation in the yeast *Saccharomyces cerevisiae*, *FEMS Microbiol. Rev.* 27 (2003) 35–64.
- [108] B.V. Geisbrecht, S.J. Gould, The human *PICD* gene encodes a cytoplasmic and peroxisomal NADP(+)-dependent isocitrate dehydrogenase, *J. Biol. Chem.* 274 (1999) 30527–30533.
- [109] B. Henke, W. Girzalsky, V. Berteaux-Lecellier, R. Erdmann, *IDP3* encodes a peroxisomal NADP-dependent isocitrate dehydrogenase required for the β -oxidation of unsaturated fatty acids, *J. Biol. Chem.* 273 (1998) 3702–3711.
- [110] C.W. van Roermund, E.H. Hetteema, A.J. Kal, M. van den Berg, H.F. Tabak, R.J. Wanders, Peroxisomal β -oxidation of polyunsaturated fatty acids in *Saccharomyces cerevisiae*: isocitrate dehydrogenase provides NADPH for reduction of double bonds at even positions, *EMBO J.* 17 (1998) 677–687.
- [111] V. Contreras-Shannon, A.P. Lin, M.T. McCammon, L. McAlister-Henn, Kinetic properties and metabolic contributions of yeast mitochondrial and cytosolic NADP⁺-specific isocitrate dehydrogenases, *J. Biol. Chem.* 280 (2005) 4469–4475.
- [112] V.D. Antonenkov, Dehydrogenases of the pentose phosphate pathway in rat liver peroxisomes, *Eur. J. Biochem.* 183 (1989) 75–82.
- [113] H. Schäfer, K. Nau, A. Sickmann, R. Erdmann, H.E. Meyer, Identification of peroxisomal membrane proteins of *Saccharomyces cerevisiae* by mass spectrometry, *Electrophoresis* 22 (2001) 2955–2968.
- [114] M. Marelli, J.J. Smith, S. Jung, E. Yi, A.I. Nesvizhskii, R.H. Christmas, R.A. Saleem, Y.Y. Tam, A. Fagarasanu, D.R. Goodlett, R. Aebersold, R. A. Rachubinski, J.D. Aitchison, Quantitative mass spectrometry reveals a role for the GTPase Rho1p in actin organization on the peroxisome membrane, *J. Cell Biol.* 167 (2004) 1099–1112.
- [115] A.G. McLennan, The Nudix hydrolase superfamily, *Cell. Mol. Life Sci.* 63 (2006) 123–143.
- [116] S.R. Abdelraheim, J.L. Cartwright, L. Gasmi, A.G. McLennan, The NADH diphosphatase encoded by the *Saccharomyces cerevisiae* NPY1 nudix hydrolase gene is located in peroxisomes, *Arch. Biochem. Biophys.* 388 (2001) 18–24.
- [117] S.R. Abdelraheim, D.G. Spiller, A.G. McLennan, Mammalian NADH diphosphatases of the Nudix family: cloning and characterization of the human peroxisomal NUDT12 protein, *Biochem. J.* 374 (2003) 329–335.
- [118] L. Gasmi, A.G. McLennan, The mouse *Nudt7* gene encodes a peroxisomal nudix hydrolase specific for coenzyme A and its derivatives, *Biochem. J.* 357 (2001) 33–38.
- [119] S.R. Abdelraheim, A.G. McLennan, The *Caenorhabditis elegans* Y87G2A.14 Nudix hydrolase is a peroxisomal coenzyme A diphosphatase, *BMC Biochem.* 3 (2002) 5.
- [120] J.L. Cartwright, L. Gasmi, D.G. Spiller, A.G. McLennan, The *Saccharomyces cerevisiae* PCD1 gene encodes a peroxisomal nudix hydrolase active toward coenzyme A and its derivatives, *J. Biol. Chem.* 275 (2000) 32925–32930.
- [121] R. Ofman, D. Spejler, R. Leen, R.J. Wanders, Proteomic analysis of mouse kidney peroxisomes: identification of RP2p as a peroxisomal nudix hydrolase with acyl-CoA diphosphatase activity, *Biochem. J.* 393 (2006) 537–543.
- [122] I. Pracharoenwattana, J.E. Cornah, S.M. Smith, *Arabidopsis* peroxisomal citrate synthase is required for fatty acid respiration and seed germination, *Plant Cell* 17 (2005) 2037–2048.
- [123] Y. Elgersma, C.W. van Roermund, R.J. Wanders, H.F. Tabak, Peroxisomal and mitochondrial carnitine acetyltransferases of *Saccharomyces cerevisiae* are encoded by a single gene, *EMBO J.* 14 (1995) 3472–3479.
- [124] L. Palmieri, F.M. Lasorsa, V. Iacobazzi, M.J. Runswick, F. Palmieri, J.E. Walker, Identification of the mitochondrial carnitine carrier in *Saccharomyces cerevisiae*, *FEBS Lett.* 462 (1999) 472–476.
- [125] C.W. van Roermund, E.H. Hetteema, M. van den Berg, H.F. Tabak, R.J. Wanders, Molecular characterization of carnitine-dependent transport of acetyl-CoA from peroxisomes to mitochondria in *Saccharomyces cerevisiae* and identification of a plasma membrane carnitine transporter, *Agp2p*, *EMBO J.* 18 (1999) 5843–5852.
- [126] I.V. Karpichev, G.M. Small, Global regulatory functions of Oaf1p and Pip2p (Oaf2p), transcription factors that regulate genes encoding peroxisomal proteins in *Saccharomyces cerevisiae*, *Mol. Cell. Biol.* 18 (1998) 6560–6570.
- [127] J.R. de Lucas, A.I. Dominguez, S. Valenciano, G. Turner, F. Laborda, The *acuH* gene of *Aspergillus nidulans*, required for growth on acetate and long-chain fatty acids, encodes a putative homologue of the mammalian carnitine/acylcarnitine carrier, *Arch. Microbiol.* 171 (1999) 386–396.
- [128] J. de Lucas, O. Martinez, P. Perez, M. Isabel Lopez, S. Valenciano, F. Laborda, The *Aspergillus nidulans* carnitine carrier encoded by the *acuH* gene is exclusively located in the mitochondria, *FEMS Microbiol. Lett.* 201 (2001) 193–198.
- [129] R.R. Ramsay, The role of the carnitine system in peroxisomal fatty acid oxidation, *Am. J. Med. Sci.* 318 (1999) 28–35.
- [130] S. Ferdinandusse, J. Mulders, S. Denis, G. Dacremont, H.R. Waterham, R.J. Wanders, Molecular cloning and expression of human carnitine octanoyltransferase: evidence for its role in the peroxisomal β -oxidation of branched-chain fatty acids, *Biochem. Biophys. Res. Commun.* 263 (1999) 213–218.
- [131] C. Indiveri, V. Iacobazzi, N. Giangregorio, F. Palmieri, Bacterial overexpression, purification, and reconstitution of the carnitine/acylcarnitine carrier from rat liver mitochondria, *Biochem. Biophys. Res. Commun.* 249 (1998) 589–594.
- [132] A.M. Lamhonwah, J. Skaug, S.W. Scherer, I. Tein, A third human carnitine/organic cation transporter (OCTN3) as a candidate for the 5q31 Crohn's disease locus (IBD5), *Biochem. Biophys. Res. Commun.* 301 (2003) 98–101.

- [133] A.M. Lamhonwah, C.A. Ackerley, A. Tilups, V.D. Edwards, R.J. Wanders, I. Tein, OCTN3 is a mammalian peroxisomal membrane carnitine transporter, *Biochem. Biophys. Res. Commun.* 338 (2005) 1966–1972.
- [134] K.A. Sacksteder, J.M. Jones, S.T. South, X. Li, Y. Liu, S.J. Gould, PEX19 binds multiple peroxisomal membrane proteins, is predominantly cytoplasmic, and is required for peroxisome membrane synthesis, *J. Cell Biol.* 148 (2000) 931–944.
- [135] F. Fraser, V.A. Zammit, Submitochondrial and subcellular distributions of the carnitine-acylcarnitine carrier, *FEBS Lett.* 445 (1999) 41–44.
- [136] S. Lawand, A.J. Dorne, D. Long, G. Coupland, R. Mache, P. Carol, *Arabidopsis* A BOUT DE SOUFFLE, which is homologous with mammalian carnitine acyl carrier, is required for postembryonic growth in the light, *Plant Cell* 14 (2002) 2161–2173.
- [137] F. Leighton, S. Bergseth, T. Rortveit, E.N. Christiansen, J. Bremer, Free acetate production by rat hepatocytes during peroxisomal fatty acid and dicarboxylic acid oxidation, *J. Biol. Chem.* 264 (1989) 10347–10350.
- [138] M.C. Hunt, S.E. Alexson, The role Acyl-CoA thioesterases play in mediating intracellular lipid metabolism, *Prog. Lipid Res.* 41 (2002) 99–130.
- [139] G. Tilton, J. Shockey, J. Browse, Two families of acyl-CoA thioesterases in *Arabidopsis*, *Biochem. Soc. Trans.* 28 (2000) 946–947.
- [140] G.B. Tilton, J.M. Shockey, J. Browse, Biochemical and molecular characterization of ACH2, an acyl-CoA thioesterase from *Arabidopsis thaliana*, *J. Biol. Chem.* 279 (2004) 7487–7494.
- [141] H. Pinfield-Wells, E.L. Rylott, A.D. Gilday, S. Graham, K. Job, T.R. Larson, I.A. Graham, Sucrose rescues seedling establishment but not germination of *Arabidopsis* mutants disrupted in peroxisomal fatty acid catabolism, *Plant J.* 43 (2005) 861–872.
- [142] K. Nicolay, M. Veenhuis, A.C. Douma, W. Harder, A 31P NMR study of the internal pH of yeast peroxisomes, *Arch. Microbiol.* 147 (1987) 37–41.
- [143] H.R. Waterham, I. Keizer-Gunnink, J.M. Goodman, W. Harder, M. Veenhuis, Immunocytochemical evidence for the acidic nature of peroxisomes in methylotrophic yeasts, *FEBS Lett.* 262 (1990) 17–19.
- [144] A.C. Douma, M. Veenhuis, G.J. Sulter, W. Harder, A proton-translocating adenosine triphosphatase is associated with the peroxisomal membrane of yeasts, *Arch. Microbiol.* 147 (1987) 42–47.
- [145] G.J. Sulter, W. Harder, M. Veenhuis, Structural and functional aspects of peroxisomal membranes in yeasts, *FEMS Microbiol. Rev.* 11 (1993) 285–296.
- [146] T.B. Dansen, K.W. Wirtz, R.J. Wanders, E.H. Pap, Peroxisomes in human fibroblasts have a basic pH, *Nat. Cell Biol.* 2 (2000) 51–53.
- [147] E.H. Pap, T.B. Dansen, K.W. Wirtz, Peptide-based targeting of fluorophores to peroxisomes in living cells, *Trends Cell Biol.* 11 (2001) 10–12.
- [148] G. Miesenböck, D.A. De Angelis, J.E. Rothman, Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins, *Nature* 394 (1998) 192–195.
- [149] A. Jankowski, J.H. Kim, R.F. Collins, R. Daneman, P. Walton, S. Grinstein, In situ measurements of the pH of mammalian peroxisomes using the fluorescent protein pHluorin, *J. Biol. Chem.* 276 (2001) 48748–48753.
- [150] T.R. van der Lende, P. Breeuwer, T. Abee, W.N. Konings, A.J. Driessen, Assessment of the microbody luminal pH in the filamentous fungus *Penicillium chrysogenum*, *Biochim. Biophys. Acta* 1589 (2002) 104–111.
- [151] I.J. van der Klei, W. Harder, M. Veenhuis, Methanol metabolism in a peroxisome-deficient mutant of *Hansenula polymorpha*: a physiological study, *Arch. Microbiol.* 156 (1991) 15–23.
- [152] T.D. Southall, S. Terhzaz, P. Cabrero, V.R. Chintapalli, J.M. Evans, J.A. Dow, S.A. Davies, Novel sub-cellular locations and functions for Secretory Pathway Ca^{2+}/Mn^{2+} ATPases, *Physiol. Genomics* 26 (2006) 35–45.
- [153] G. Wu, A.J. Robertson, P. Zheng, X. Liu, L.V. Gusta, Identification and immunogold localization of a novel bromegrass (*Bromus inermis* Leyss) peroxisome channel protein induced by ABA, cold and drought stresses, and late embryogenesis, *Gene* 363 (2005) 77–84.
- [154] Y. Elgersma, H.F. Tabak, Proteins involved in peroxisome biogenesis and functioning, *Biochim. Biophys. Acta* 1286 (1996) 269–283.